

GRADY'S 2024 STROKE QUALITY MEASURES

(Source: Get With The Guidelines)

Metric	Description	Our Score	National Average Score (all hospitals)
Anticoagulation for Afib/Aflutter	Percentage of patients with an ischemic stroke or TIA with atrial fibrillation/flutter discharged on anticoagulation therapy	100%	97%
Anti-thrombotics at Discharge	Percentage of patients with an ischemic stroke or TIA prescribed antithrombotic therapy at discharge.	99%	98%
Intensive Statin Therapy	Percentage of Ischemic Stroke and TIA patients who are prescribed high-intensity statin therapy at discharge OR, if > 75 years of age, are prescribed at least moderate intensity statin therapy at discharge	100%	95%
Stroke Education	Percentage of patients with Stroke or TA or their caregivers who were given education and/or educational materials during the hospital stay addressing ALL of the following: personal risk factors for stroke, warning signs for stroke, activation of emergency medical system, need for follow-up after discharge, and medications prescribed.	99%	94%
Smoking Cessation	Percentage of patients with ischemic or hemorrhagic stroke, or TIA with a history of smoking cigarettes, who are, or whose caregivers are, given smoking cessation advice or counseling during hospital stay.	100%	98%
VTE Prophylaxis	Percentage of patients with an ischemic stroke, or a hemorrhagic stroke, or stroke not otherwise specified who receive VTE prophylaxis the day of or the day after hospital admission	97%	96%
Rehabilitation Considered	Percentage of patients with Stroke who were assessed for rehabilitation services during hospital stay.	100%	99%
Dysphagia Screening	Percentage of Stroke patients who undergo screening for dysphagia with an evidence based bedside testing protocol approved by the hospital before being given any food, fluids, or medication by mouth.	92%	84%
Procoagulant Reversal	Percentage of Procoagulant Reversal Agent Initiation for Intracerebral Hemorrhage (ICH).	100%	92%
IV Fibrinolytic (Tenecteplase)			
IV Tenecteplase Arrive by 2 Hour, Treat by 3 Hour	Percentage of acute ischemic stroke patients who arrive at the hospital within 120 minutes (2 hours) of time last known per hospital for whom IV thrombolytic was initiated at this hospital within 180 minutes (3 hours) of time last known well.	100%	92%
IV Alteplase Arrive by 3.5 Hour, Treat by 4.5 Hour	Percentage of acute ischemic stroke patients who arrive at the hospital within 210 minutes (3.5 hours) of time last known well and for whom IV thrombolytic was initiated at this hospital within 270 minutes (4.5 hours) of time last known well.	100%	94%
Thrombectomy			
Thrombolysis in Cerebral Infarction (TICI Post-Treatment Reperfusion Score - TICI 2B or greater)	Ischemic stroke patients with a post-treatment reperfusion grade of TICI 2B or higher in the vascular territory beyond the target arterial occlusion at the end of treatment with intra-arterial (IA) thrombolytic (tPA/TNK) therapy and/or mechanical endovascular reperfusion therapy	95%	89%
Rate of Rapid Effective Reperfusion from Hospital Arrival	Ischemic Stroke Patients with a large vessel cerebral occlusion who receive mechanical endovascular reperfusion (MER) therapy within 120 minutes of patient arrival	84%	72%